



NORTH-HOLLAND

On $(A, B)^t$ -Invariant Subspaces Having Extendible Brunovsky Bases

A. Compta* and J. Ferrer†

*Departament de Matemàtica Aplicada I
E.T.S. Enginyers Industrials de Barcelona
Universitat Politècnica de Catalunya
Diagonal 647
08028 Barcelona, Spain*

Submitted by Leiba Rodman

ABSTRACT

We consider $(A, B)^t$ -invariant subspaces having a Brunovsky basis which can be extended to a Brunovsky basis of the whole space. We obtain a geometrical characterization of this class of $(A, B)^t$ -invariant subspaces, and a complete family of numerical invariants to classify them. © Elsevier Science Inc., 1997

1. INTRODUCTION

For A a square matrix, Gohberg et al. [6] introduce an “interesting class” of A -invariant subspaces, which they call “marked”: an A -invariant subspace is marked if and only if it has a Jordan basis which can be extended to a Jordan basis of the space. The A -marked subspaces have been studied, for example, in [2] and [5].

Moreover, for (A, B) a pair of matrices, [6], [3] and other works define and study the (A, B) -invariant subspaces and the $(A, B)^t$ -invariant subspaces.

*E-mail: compta@ma1.upc.es.

†We are very grateful to F. Puerta and X. Puerta for their valuable comments and suggestions. We also thank the referees for their careful review.

The question naturally arises of the existence of Brunovsky bases of those subspaces which can be extended to Brunovsky bases of the space. (A basis is called a *Brunovsky basis* if relative to it the matrix has the Brunovsky reduced form.) That is to say, we wish to consider (A, B) -marked and $(A, B)^t$ -marked subspaces.

This paper is a geometrical approach to the study of $(A, B)^t$ -marked subspaces. In fact, following the techniques in [4], we consider linear maps $f: Y \rightarrow X$ defined only on a subspace of Y of X . Then we say that a subspace $W \subset Y$ is f -invariant if $f(W) \cap Y \subset W$, and among these subspaces we call f -marked those having a extendible Brunovsky basis.

The first goal is a geometrical characterization of the f -marked subspaces. In [5], such a characterization has been obtained for f an endomorphism. Here, Theorem 5.1 generalizes this result by means of the additional condition

$$[W \cap f^{j-i}(f^{-j+1}(Y))] + Y_i = (W + Y_i) \cap [f^{j-i}(f^{-j+1}(Y)) + Y_i]$$

for all $1 \leq i \leq j$.

The proof of this theorem includes the construction of a Brunovsky basis of W and its extension to a Brunovsky basis of $f: Y \rightarrow X$, provided that the above conditions are verified. This explicit construction suggests a condition to guarantee the existence of a bijection between these kind of bases for two different f -marked subspaces. Thus, the second main result is Theorem 6.1, which classifies the $(A, B)^t$ -marked subspaces by means of a complete family of numerical invariants.

Section 2 contains the notation which will be used in the paper.

In Sections 3 and 4 we present the definition and some generalities of $(A, B)^t$ -invariant and $(A, B)^t$ -marked subspaces respectively. In particular, in Section 3.5 we characterize the Brunovsky invariants of an f -invariant subspace. This result has been proved in [3] by means of matricial techniques.

Section 5 is devoted to proving Theorem 5.1.

In Section 6 we define a natural equivalence between $(A, B)^t$ -invariant subspaces, and we prove the classification Theorem 6.1. As an application, in Section 6.7 we compute the number of nonequivalent f -marked subspaces for a fixed linear map $f: Y \rightarrow X$.

2. PRELIMINARIES

2.1

X will be an $(n + m)$ -dimensional vector space over the complex numbers \mathbb{C} .

If B is a subset of X , the symbol $[B]$ will denote the subspace spanned by the vectors of B .

If $X \supset E_1 \supset E_2 \supset \dots$ is a chain of subspaces of X , we say that a basis B of X is *adapted* to it if $B \cap E_i$ is a basis of E_i for all $i = 1, 2, \dots$.

2.2

$Y \subset X$ will be an n -dimensional subspace, and $f: Y \rightarrow X$ a linear map defined on it. In any basis of X adapted to $Y \subset X$, the matrix f is a pair

$$\begin{pmatrix} A \\ B \end{pmatrix} \quad \text{where} \quad A \in M_n(\mathbb{C}) \quad \text{and} \quad B \in M_{m,n}(\mathbb{C}).$$

As usual, we will say that two linear maps defined on a subspace, $f: Y \rightarrow X$ and $f': Y' \rightarrow X'$, are equivalent (or *block-similar*) if there is an isomorphism $\varphi: X \rightarrow X'$ such that $\varphi(Y) = Y'$, $\varphi \circ f = f' \circ \hat{\varphi}$, where $\hat{\varphi}$ is the restriction of φ to Y . It amounts to saying that their respective matrices

$$\begin{pmatrix} A \\ B \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} A' \\ B' \end{pmatrix}$$

are block-similar.

2.3

Following [4], we will consider the stationary chain of subspaces and linear maps defined as follows:

$$Y_0 \equiv Y, \quad Y_i = f^{-1}(Y_{i-1}), \quad i > 0,$$

$$Y_{k+1} = Y_k \subset Y_{k-1} \subset \dots \subset Y_1 \subset Y \subset X,$$

$$f_i: Y_i \rightarrow Y_{i-1}, \quad i > 0,$$

where f_i is the corresponding restriction of f . We will denote f_i simply by f if no confusion is possible.

2.4

We recall that a complete family of invariants of the equivalence in Section 2.2 is formed by:

(1) The Kronecker indices (k_1, \dots, k_r) , which can be computed as the conjugate partition of (r_1, \dots, r_k) , where

$$r_i = \dim Y_{i-1} - \dim Y_i, \quad i = 1, \dots, k.$$

(2) The similarity invariants of the endomorphism

$$f_{k+1} : Y_{k+1} \rightarrow Y_k$$

or equivalently its invariant factors $\alpha_1 \mid \alpha_2 \mid \dots \mid \alpha_t$.

In particular, $f : Y \rightarrow X$ is observable if and only if $Y_k = \{0\}$.

2.5

Also, we recall that a Brunovsky basis of $f : Y \rightarrow X$ is formed by a Jordan basis of $f_{k+1} : Y_{k+1} \rightarrow Y_k$ joint with r so-called *Brunovsky chains*

$$x_j, f(x_j), \dots, f^{k_i}(x_j) \notin Y, \quad 1 \leq j \leq r,$$

such that $f^{k_1}(x_1), \dots, f^{k_r}(x_r)$ are linearly independent. They can be obtained by taking a Jordan basis of Y_k and extending it successively in the chain of subspaces $\dots \subset Y_i \subset \dots$ by taking images and extending step by step—that is to say, by means of supplementary subspaces \bar{Y}_i ($i = k, \dots, 1, 0$) such that

$$Y_{k-1} = Y_k \oplus \bar{Y}_k,$$

$$Y_{k-2} = Y_{k-1} \oplus \bar{Y}_{k-1}, \quad \bar{Y}_{k-1} \supset f(\bar{Y}_k),$$

$$\vdots$$

$$Y = Y_1 \oplus \bar{Y}_1, \quad \bar{Y}_1 \supset f(\bar{Y}_2),$$

$$X = Y \oplus \bar{Y}, \quad \bar{Y} \supset f(\bar{Y}_1).$$

3. INVARIANT SUBSPACES UNDER A LINEAR MAP DEFINED ON A SUBSPACE

3.1

According to [6] [Theorem 6.6.1, with $X = \mathbb{C}^n \times \mathbb{C}^m$ and $Y = \mathbb{C}^n \equiv \mathbb{C}^n \times \{0\} \subset \mathbb{C}^n \times \mathbb{C}^m$, so that $\ker C = f^{-1}(Y)$] and [3], we define:

DEFINITION 3.1. Let $f: Y \rightarrow X$ be a linear map defined on a subspace, and $W \subset Y$ a subspace. We say that W is f -invariant if

$$f(W) \cap Y \subset W.$$

3.2. Example

Let $f: Y \rightarrow X$ be a linear map defined on a subspace. If $y, f(y), \dots, f^k(y)$ is a full Brunovsky chain, then the subspaces generated by subchains of the kind $f^d(y), f^{d+1}(y), \dots, f^{k-1}(y)$ are f -invariant. We will see in the next section that these subspaces are in fact “ f -marked” subspaces.

3.3

Then we have the following matricial characterization:

PROPOSITION 3.1. Let $f: Y \rightarrow X$ and $W \subset Y$ be as in Section 3.1. Then W is f -invariant if and only if there is a basis of X adapted to $W \subset Y \subset X$ such that the matrix $\begin{pmatrix} A \\ B \end{pmatrix}$ of f in this basis has the form

$$\begin{pmatrix} A_1^1 & A_2^1 \\ 0 & A_2^2 \\ B_1 & B_2 \end{pmatrix}$$

where $A_s^1 \in M_s(\mathbb{C})$, $s = \dim W$.

Proof. Obviously, Definition 3.1 is the same as saying that there is a subspace $\bar{Y} \subset X$ such that

$$Y \oplus \bar{Y} = X, \quad f(W) \subset W \oplus \bar{Y}.$$

Then, in any basis of X adapted to $W \subset Y \subset X$ and also to $\bar{Y} \subset X$, the matrix of f has the desired form. The converse is obvious. ■

3.4

For a geometrical approach, we will consider the restriction of f to W ,

$$\hat{f}: W \rightarrow X,$$

as a linear map defined on a subspace, and, in an analogous way to (2.3), the chain of subspaces and linear maps

$$W_0 \equiv W, \quad W_i = \hat{f}^{-1}(W_{i-1}), \quad i > 0,$$

$$W_{h+1} = W_h \subset W_{h-1} \subset \cdots \subset W_1 \subset W \subset X,$$

$$\hat{f}_i : W_i \rightarrow W_{i-1}, \quad i > 0.$$

Obviously, we have

$$\begin{array}{ccccccccccccccc} \cdots & \subset & Y_i & \subset & Y_{i-1} & \subset & \cdots & \subset & Y_1 & \subset & Y & \subset & X \\ & & \cup & & \cup & & & & \cup & & \cup & & \\ \cdots & \subset & W_i & \subset & W_{i-1} & \subset & \cdots & \subset & W_1 & \subset & W & \subset & X \end{array}$$

3.5

Then we have the following characterization:

PROPOSITION 3.2. *Let $f : Y \rightarrow X$ be a linear map defined on a subspace, and $W \subset Y$ a subspace. Then, with the above notation, the following statements are equivalent:*

- (i) W is f -invariant.
- (ii) $W_1 = W \cap Y_1$.
- (iii) $W_i = W_{i-1} \cap Y_i = W \cap Y_i$ for all $i > 0$.

Proof. The equivalence (i) \Leftrightarrow (ii) is straightforward. And (iii) can be easily proved by induction:

$$\begin{aligned} W_{i+1} &= \hat{f}^{-1}(W_i) = \hat{f}^{-1}(W \cap Y_i) = \hat{f}^{-1}(W \cap Y_i) \cap W \\ &= \hat{f}^{-1}(W) \cap Y_{i+1} \cap W = \hat{f}^{-1}(W) \cap Y_{i+1} = W_1 \cap Y_{i+1} \\ &= W \cap Y_i \cap Y_{i+1} = W \cap Y_{i+1}. \end{aligned} \quad \blacksquare$$

3.6

This geometric approach allows an alternative proof to the relation in [3] between the Brunovsky invariants of $f : Y \rightarrow X$ and those of $\hat{f} : W \rightarrow X$.

PROPOSITION 3.3 [3]. *Let $f: Y \rightarrow X$ be a linear map defined on a subspace, and $W \subset Y$ an f -invariant subspace. We denote the Brunovsky invariants of $f: Y \rightarrow X$, as in (2.4), and in an analogous way we denote by $(\hat{k}_1, \hat{k}_2, \dots, \hat{k}_q)$ and $\hat{\alpha}_1 | \hat{\alpha}_2 | \dots | \hat{\alpha}_l$ those of $\hat{f}: W \rightarrow X$. Then, we have*

- (i) $l \leq t$ and $\alpha_i | \hat{\alpha}_i | \alpha_{i+t-l}$ for $i = 1, 2, \dots, l$;
- (ii) $q \leq r$ and $\hat{k}_i \leq k_i$ for $i = 1, 2, \dots, q$.

Conversely, given $(\hat{k}_1, \dots, \hat{k}_q)$ and $\hat{\alpha}_1 | \hat{\alpha}_2 | \dots | \hat{\alpha}_l$ verifying (i) and (ii), there is an f -invariant subspace $W \subset Y$ such that they are the Brunovsky invariants of $\hat{f}: W \rightarrow X$.

Proof. Obviously $k \geq h$, and the subspace $W_h = W_k \subset Y_k$ is invariant under the endomorphism $f_{k+1}: Y_k \rightarrow Y_k$. Then (i) is well known [1, p. 149]. To proof (ii), we consider the conjugate partitions (r_1, \dots, r_k) and $(\hat{r}_1, \dots, \hat{r}_h)$ of (k_1, \dots, k_r) and $(\hat{k}_1, \dots, \hat{k}_q)$ respectively (see Section 2.4), where

$$r_i = \dim Y_{i-1} - \dim Y_i,$$

$$\hat{r}_i = \dim W_{i-1} - \dim W_i.$$

From (iii) of Proposition 3.2 we obtain $r_i \geq \hat{r}_i$. That obviously implies $k_i \geq \hat{k}_i$ and $r \geq q$. For the converse, it is sufficient to take as W the subspace spanned by a set of Brunovsky subchains having the desired length (see Section 3.2). ■

3.7

As we have said in Section 3.2, we will see in the next section that the f -invariant subspace constructed in the converse of the above proof is in fact a so-called f -marked subspace. In Section 6.7, for f observable, we will determine the number of "different" subspaces of this kind for each $(\hat{k}_1, \dots, \hat{k}_q)$ verifying relation (ii).

4. MARKED SUBSPACES UNDER A LINEAR MAP DEFINED ON A SUBSPACE

4.1

In all this section, we assume $f: Y \rightarrow X$ a linear map defined in a subspace, $W \subset Y$ a f -invariant subspace, and $\hat{f}: W \rightarrow X$ the restriction of f . We maintain the notation in Sections 2 and 3.4.

4.2

Gohberg et al. [6] define the *marked* invariant subspaces under an endomorphism by means of the following condition: there is a Jordan basis of the subspace which can be extended to a Jordan basis of the global space. In an analogous way, we define:

DEFINITION 4.1. In the conditions of Section 4.1, we say that W is f -marked if there is a Brunovsky basis of $\hat{f}: W \rightarrow X$ which can be extended to a Brunovsky basis of $f: Y \rightarrow X$, or equivalently, if there is a Brunovsky basis of $f: Y \rightarrow X$ adapted to $W \subset Y \subset X$.

4.3

In a similar way to Section 3.3, the following matricial characterization is immediate:

PROPOSITION 4.1. *With the above notation, W is f -marked if and only if there is a basis of X adapted to $W \subset Y \subset X$ such that the matrix $\begin{pmatrix} A \\ B \end{pmatrix}$ of f in this basis has the form*

$$\begin{pmatrix} A_1^1 & A_2^1 \\ 0 & A_2^2 \\ B_1 & B_2 \end{pmatrix}, \quad \text{where } A_1^1 \in M_s(\mathbb{C}), \quad s = \dim W,$$

and in addition:

(i) $\begin{pmatrix} A_1^1 \\ B_1 \end{pmatrix}$ is a Brunovsky matrix.

(ii) $\begin{pmatrix} A \\ B \end{pmatrix}$ is a Brunovsky matrix up to permutation; that is to say, there is a permutation matrix $P \in \text{GL}(n, \mathbb{C})$ such that

$$\begin{pmatrix} P^t & 0 \\ 0 & \text{Id} \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix} P$$

is a Brunovsky matrix.

4.4. Example

Let $X = \mathbb{C}^4$, (e_1, e_2, e_3, e_4) be a basis, $Y = [e_1, e_2, e_3]$, and $f: Y \rightarrow X$ be defined by

$$f(e_1) = 0, \quad f(e_2) = e_1 + e_3, \quad f(e_3) = e_4.$$

Then $W = [e_3]$ is f -invariant, but it is not f -marked.

5. GEOMETRIC CHARACTERIZATION OF f -MARKED SUBSPACES

5.1

In [5] the authors characterize the f -marked subspaces for $f: X \rightarrow X$ an endomorphism. In fact, for f nilpotent, they prove that an invariant subspace $W \subset X$ is marked if and only if

$$\begin{aligned} W \cap f^{d+1}(K_{d+h+1}) + W \cap f^d(K_{d+h-1}) \\ = W \cap [f^{d+1}(K_{d+h+1}) + f^d(K_{d+h-1})] \end{aligned}$$

for all d, h , where $K_l = \ker f^l = f^{-l}(0)$.

Here, for $f: Y \rightarrow X$ a linear map defined on a subspace, we obtain a similar geometric characterization of the f -marked subspaces $W \subset Y$ in terms of the subspaces $Y_i = f^{-i}(Y)$ (see Section 2.3) and $W_i = \hat{f}^{-i}(W)$ (see Section 3.4). It is proved in Sections 5.3, 5.4, and 5.5.

THEOREM 5.1. *Let $f: Y \rightarrow X$ be a linear map defined on a subspace, and $W \subset Y$ an f -invariant subspace. Then W is f -marked if and only if*

- (i) W_k is f_{k+1} -marked,
- (ii) $[W \cap f^{j-i}(Y_{j-1})] + Y_i = (W + Y_i) \cap [f^{j-i}(Y_{j-1}) + Y_i]$

for all $1 \leq i \leq j \leq k$.

5.2. Remarks

REMARK 1. Because $f^{j-i}(Y_{j-1}) \subset Y_{i-1}$ and $W \cap Y_{i-1} = W_{i-1}$ (see Section 3.5), we can replace W by W_{i-1} in both members of (ii). Furthermore, the right inclusion in (ii) is always true.

Hence, condition (ii) above is equivalent to

$$(ii') [W_{i-1} \cap f^{j-i}(Y_{j-1})] + Y_i \supset (W_{i-1} + Y_i) \cap [f^{j-i}(Y_{j-1}) + Y_i].$$

REMARK 2. Notice that this inclusion (ii') is always true for $i = j$, and for $i > h$.

REMARK 3. X. Puerta has remarked that condition (ii) is equivalent to

$$(ii'') W \cap [f^{j-i}(Y_{j-1}) + Y_i] = W \cap f^{j-i}(Y_{j-1}) + W \cap Y_i,$$

which is much more similar to the one for f a nilpotent endomorphism.

5.3. Proof of the Necessity

(i) is obvious.

(ii): With the notation in Section 2.5, we have, for all $1 \leq i \leq j \leq k$,

$$f^{j-i}(Y_{j-1}) + Y_i = [f^{j-i}(Y_j) \oplus f^{j-i}(\bar{Y}_j)] + Y_i = f^{j-i}(\bar{Y}_j) + Y_i$$

(in fact, the last sum is a direct sum). The Brunovsky bases of $\hat{f}: W \rightarrow X$ are obtained in an analogous way to (2.5). In particular, for all $1 \leq i \leq k$,

$$W_{i-1} + Y_i = \bar{W}_i + Y_i$$

(here also, the last sum is a direct sum).

Since W is f -marked, there exist Brunovsky bases of $f: Y \rightarrow X$ and $\hat{f}: W \rightarrow X$ such that

$$\bar{W}_i = \bar{Y}_i \cap W \quad (1 \leq i \leq k).$$

Hence, for all $1 \leq i < j \leq k$,

$$[\bar{W}_i + f^{j-i}(\bar{Y}_j)] \cap Y_i = 0.$$

Therefore

$$(\bar{W}_i + Y_i) \cap [f^{j-i}(\bar{Y}_j) + Y_i] = [\bar{W}_i \cap f^{j-i}(\bar{Y}_j)] + Y_i,$$

and inclusion (ii') in Section 5.2 follows immediately.

5.4

To prove the sufficiency, we will need the following lemma:

LEMMA 5.1. *In the conditions of Section 4.1, let*

$$W_j^i = \frac{(W_{i-1} + Y_i) \cap [f^{j-i}(Y_{j-1}) + Y_i]}{(W_{i-1} + Y_i) \cap [f^{j-i+1}(Y_j) + Y_i]}, \quad 1 \leq i \leq j \leq k.$$

Then

(a) We have natural isomorphisms

$$\frac{W}{W_k} \cong \bigoplus_{i=1}^k \frac{W_{i-1}}{W_i},$$

$$\frac{W_{i-1}}{W_i} \cong \bigoplus_{j=i}^k W_j^i \quad (1 \leq i \leq k).$$

(b) For all $1 \leq i \leq j \leq k$, the linear maps

$$\tilde{f}: W_j^i \rightarrow W_j^{i-1}$$

induced by f are injective.

Proof. (a): The first isomorphism is obvious, bearing in mind the stationary chain

$$\cdots = W_k = \cdots = W_h \subset W_{h-1} \subset \cdots \subset W_1 \subset W.$$

For the second one, we remark that, because W is f -invariant, we have (see Section 3.7) $W_i = W_{i-1} \cap Y_i$, so that

$$\frac{W_{i-1}}{W_i} = \frac{W_{i-1}}{W_{i-1} \cap Y_i} \cong \frac{W_{i-1} + Y_i}{Y_i}.$$

Now, we consider the finite chain

$$Y_i = f^{k+1-i}(Y_k) + Y_i \subset \cdots \subset f^{j-i}(Y_{j-1}) + Y_i \subset \cdots \subset Y_{i-1} + Y_i = Y_{i-1}.$$

If we intersect each term with $W_{i-1} + Y_i$, we have

$$Y_i \subset \cdots \subset (W_{i-1} + Y_i) \cap [f^{j-i}(Y_{j-1}) + Y_i] \subset \cdots \subset W_{i-1} + Y_i,$$

and the desired isomorphism follows immediately.

(b): Let $x \in f^{j-i}(Y_{j-1}) + Y_i$, with $\tilde{f}(\tilde{x}) = 0$. Then there exists $y \in Y_j$ such that $f(x) - f^{j-i+2}(y) \in Y_{i-1}$. Hence $x - f^{j-i+1}(y) \in Y_i$, and $\tilde{x} = 0$. ■

5.5. Proof of the Sufficiency

Firstly, because of condition (i) we can obtain a Jordan basis of Y_k adapted to W_k . Hence it is sufficient to construct a Brunovsky basis of a complementary subspace of W_k to W which can be extended to a Brunovsky basis of a complementary subspace of Y_k to Y . In order to do so, we can sketch the above lemma in the following diagram

$$\begin{array}{c}
 W/W_k \\
 \cong \\
 W_{k-1}/W_k \cong W_k^k \\
 \oplus \searrow \\
 W_{k-2}/W_{k-1} \cong W_{k-1}^{k-1} \oplus W_k^{k-1} \\
 \oplus \searrow \searrow \\
 W_{k-3}/W_{k-2} \cong W_{k-2}^{k-2} \oplus W_{k-1}^{k-2} \oplus W_k^{k-2} \\
 \oplus \vdots \searrow \\
 \vdots \\
 \oplus \\
 W/W_1 \cong W_1^1 \oplus W_2^1 \oplus \cdots \oplus W_k^1
 \end{array}$$

But now, if condition (ii) in Section 5.1 (or equivalently, condition (ii') in Section 5.2) is verified, we have, for all $1 \leq i \leq j \leq k$

$$W_j^i = \frac{[W_{i-1} \cap f^{j-i}(Y_{j-1})] + Y_i}{[W_{i-1} \cap f^{j-i+1}(Y_j)] + Y_i} \cong \frac{W_{i-1} \cap f^{j-i}(Y_{j-1})}{W_{i-1} \cap f^{j-i+1}(Y_j) + W_i \cap f^{j-i}(Y_{j-1})}$$

(the last step is a direct application of the general natural isomorphism

$$\frac{F + G}{F' + G} \cong \frac{F}{F' + F \cap G},$$

where $F' \subset F$).

Thus, we can obtain an extendible Brunovsky basis of W by means of taking images and extending step by step (see Section 2.5), as we sketch in the following diagram:

$$\begin{array}{ccccccc}
 \hat{B}_k^k & & & & & & \\
 \hat{B}_{k-1}^{k-1} & f(\hat{B}_k^k), \hat{B}_k^{k-1} & & & & & \\
 \hat{B}_{k-2}^{k-2} & f(\hat{B}_{k-1}^{k-1}), \hat{B}_{k-1}^{k-2} & f^2(\hat{B}_k^k), f(\hat{B}_k^{k-1}), \hat{B}_k^{k-2} & & & & \\
 \vdots & \vdots & \vdots & & & & \\
 \hat{B}_1^1 & f(\hat{B}_2^2), \hat{B}_1^1 & \cdots & \cdots & f^{k-1}(\hat{B}_k^k), \dots, f(\hat{B}_k^2), \hat{B}_k^1 & &
 \end{array}$$

That is to say, we take:

for $j = k, k-1, \dots, 1$: $\hat{B}_j^j \subset W_{j-1} \cap Y_{j-1}$ such that its classes form a basis of W_j^j ,

for $j = k, k-1, \dots, 2$: $\hat{B}_j^{j-1} \subset W_{j-2} \cap f(Y_{j-1})$ such that the classes of $f(\hat{B}_j^j), \hat{B}_j^{j-1}$ form a basis of W_j^{j-1} ,

...

for $j = k$: $\hat{B}_j^{j-k+1} \subset W_{j-k} \cap f^{k-1}(Y_{j-1})$ such that the classes of $f^{k-1}(\hat{B}_j^j), \dots, f(\hat{B}_j^{j-k+2}), \hat{B}_j^{j-k+1}$ form a basis of W_j^{j-k+1} .

Obviously, the above sets \hat{B}_j^i , $1 \leq i \leq j \leq k$, and their images form a Brunovsky basis \hat{B} of the complementary subspace $\bigoplus_{i=1}^k \bar{W}_i$ of W_k to W where

$$\bar{W}_i = \left[\bigcup_{\substack{i \leq j \leq k \\ 0 \leq l \leq j-i}} f^l(\hat{B}_j^{i+l}) \right], \quad 1 \leq i \leq k.$$

And this basis is extendible to a Brunovsky basis of a complementary subspace of Y_k to Y because

$$\hat{B}_j^i \subset W_{i-1} \cap f^{j-i}(Y_{j-1}), \quad 1 \leq i \leq j \leq k.$$

In fact, first we extend the chains in \hat{B} to their maximal length, taking antiimages

$$f^{-1}(\hat{B}_j^i), \dots, f^{-(j-i)}(\hat{B}_j^i), \quad 1 \leq i < j \leq k,$$

where this doesn't mean the set of antiimages of \hat{B}_j^i , but a chain of antiimages of each vector in \hat{B}_j^i .

Finally, if we write

$$\hat{Y}_i = \bigoplus_{j \geq i}^k \left[f^{j-i}(\hat{B}_j^j), f^{j-i-1}(\hat{B}_j^{j-1}), \dots, \hat{B}_j^i, f^{-1}(\hat{B}_j^{i-1}), \dots, f^{-i+1}(\hat{B}_j^1) \right]$$

for $1 \leq i \leq k$, we have

$$Y_i \oplus \bar{W}_i \subset Y_i \oplus \hat{Y}_i \subset Y_{i-1},$$

$$f(\hat{Y}_i) \subset \hat{Y}_{i-1}.$$

Therefore, we can construct $\bar{Y}_k, \bar{Y}_{k-1}, \dots$ (see Section 2.5), such that $\bar{Y}_i \supset \hat{Y}_i$. Thus, we have extended the basis B of $\bigoplus_{k=1}^k \bar{W}_i$ to a Brunovsky basis of the complementary subspace $\bigoplus_{i=1}^k \bar{Y}_i$ of Y_k to Y , as we desired.

6. CLASSIFICATION OF f -MARKED SUBSPACES

6.1

We consider the following natural equivalence relation between two f -marked subspaces:

DEFINITION 6.1. Let $f: Y \rightarrow X$, $f': Y' \rightarrow X'$ be two linear maps defined on a subspace, and $W \subset Y$, $W' \subset Y'$ marked subspaces under them respectively. We say that they are equivalent if there is an isomorphism $\varphi: X \rightarrow X'$ such that

$$\varphi(Y) = Y', \quad \varphi(W) = W',$$

$$\varphi \circ f = f' \circ \hat{\varphi},$$

where $\hat{\varphi}$ is the restriction of φ to Y .

6.2

In particular, f and \hat{f} must be block-similar to f' and \hat{f}' respectively. Obviously, we must suppose $\dim X = \dim X'$, $\dim Y = \dim Y'$, $\dim W = \dim W'$. From now on, these dimensions will be respectively $n + m$, n , and s .

6.3

The construction in Sections 5.4 and 5.5 suggests considering the invariants $\dim W_j^i$:

DEFINITION 6.2. Let $f: Y \rightarrow X$ be a linear map defined on a subspace, and $W \subset Y$ an f -marked subspace. We consider the numerical invariants

$$\begin{aligned} w_j^i &= \dim (W + Y_i) \cap [f^{j-i}(Y_{j-1}) + Y_i] \\ &\quad - \dim (W + Y_i) \cap [f^{j-i+1}(Y_j) + Y_i] \end{aligned}$$

for $1 \leq i \leq j \leq k$. For convenience, we define $w_j^i = 0$ if $1 \leq j < i$.

6.4

From (a) in Lemma 5.1 it is obvious that $\hat{r}_i = \dim W_{i-1} - \dim W_i = w_i^i + \dots + w_k^i$ ($1 \leq i \leq k$), and we recall that the conjugate partition of $(\hat{r}_1, \dots, \hat{r}_k)$ gives the Kronecker indices of $\hat{f}: W \rightarrow X$.

6.5

Then we have the following classification result.

THEOREM 6.1. *Let $f: Y \rightarrow X$, $f': Y' \rightarrow X'$ be two linear maps defined on a subspace, and $W \subset Y$, $W' \subset Y'$ marked subspaces under them respectively. Then they are equivalent if and only if:*

- (i) *f and f' are block-similar.*
- (ii) *W_k and W'_k are equivalent as marked subspaces under the endomorphisms f_{k+1} and f'_{k+1} , respectively.*
- (iii) *$w_j^i = w_j'^i$ for all $1 \leq i \leq j \leq k$.*

Proof. The three conditions are obviously necessary. For the sufficiency, we will consider the construction of extendible Brunovsky bases in Section 5.5, to show that a bijection between Brunovsky bases of X and X' , adapted to $W \subset Y \subset X$ and $W' \subset Y' \subset X'$ respectively, and commuting with f and f' , is possible. Because of condition (ii), this is trivial for the Jordan bases of Y_k and Y'_k . Moreover, condition (iii) implies that the numbers of vectors in the sets \hat{B}_j^i , which generate the Brunovsky chains for \hat{f} , are the same as the numbers of the analogous ones for \hat{f}' ; then, if we take bijections $\hat{B}_j^i \rightarrow \hat{B}_j'^i$, $1 \leq i \leq j \leq k$, and we extend these bijections naturally to the corresponding generated chains, we obtain a bijection $\hat{B} \rightarrow \hat{B}'$ which commutes with f and f' . Now, we extend the bijection to the antiimages $f^{-1}(\hat{B}_j^i)$, so that we have isomorphisms $\hat{Y}_i \rightarrow \hat{Y}'_i$, $1 \leq i \leq k$, always commuting with f and f' . Finally, condition (i) ensures that \bar{Y}_i, \bar{Y}'_i can be chosen so that the above isomorphisms can be extended to $\bar{Y}_i \rightarrow \bar{Y}'_i$, $1 \leq i \leq k$, commuting with f and f' . ■

6.6

We remark that [5] gives conditions for (ii) above to be verified.

6.7

As an application, we can determine how many nonequivalent marked subspaces verify the converse in Proposition 3.3 for f observable.

PROPOSITION 6.1. *Let $f: Y \rightarrow X$ be an observable linear map defined on a subspace, with Kronecker indices $k_1 \geq \dots \geq k_r$. Then there are as many (up to equivalence) f -marked subspaces $W \subset Y$ as collections of integers (m_1, \dots, m_r) verifying the following conditions:*

- (i) $k_i \geq m_i \geq 0$;
- (ii) $m_i \geq m_{i+1}$ for the indices $1 \leq i \leq r$ such that $k_i = k_{i+1}$.

In particular, given $\hat{k}_1 \geq \dots \geq \hat{k}_r$ with $\hat{k}_i \leq k_i$ ($1 \leq i \leq r$), there are as many (up to equivalence) f -marked subspaces $W \subset Y$ with Kronecker indices $(\hat{k}_1, \dots, \hat{k}_r)$ as permutations (m_1, \dots, m_r) of them verifying (i) and (ii).

Proof. In Theorem 6.1 we saw that an f -marked subspace is characterized up to equivalence by the numbers w_j^i , $1 \leq i \leq j \leq k$. These numbers must verify

- (a) $w_j^i \geq w_j^{i+1} \geq 0$,
- (b) $w_j^1 \leq r_j - r_{j+1}$,

and conversely, it is obvious that for every family of numbers w_j^i , $1 \leq i \leq j \leq k$, verifying (a) and (b) there exists an f -marked subspace which has these invariants. Therefore, we have proved that there is a bijection between the sets of numbers $\{w_j^i\}$ verifying conditions (a)–(b) and the equivalence classes of f -marked subspaces.

To conclude, we will prove that there exists a bijection between the collections of numbers $\{w_j^i\}$ verifying (a)–(b) and those of numbers $\{m_i\}$ verifying conditions (i)–(ii) in the proposition.

Given $\{w_j^i\}_{1 \leq i \leq j \leq k}$ with conditions (a)–(b), if $r_{j+1} < i \leq r_j$, we define $m_i = \text{card}_{1 \leq l \leq j} \{w_j^l : w_j^l \geq i - r_{j+1}\}$, and if $w_j^1 > 0$, then $m_{r_{j+1}+1} \geq \dots \geq m_{r_j}$ is the conjugated partition of $w_j^1 \geq \dots \geq w_j^j$ filled by zeros if $w_j^1 < r_j - r_{j+1}$.

Let us see that this collection verifies conditions (i)–(ii) in the proposition.

(i): If $r_{j+1} < i \leq r_j$, then obviously $m_i \leq j$. But, because of the relation between k -numbers and r -numbers, we have $k_{r_{j+1}+1} = \dots = k_{r_j} = j$.

(ii): If $k_i = k_{i+1}$, there exists j ($1 \leq j \leq k$) such that $r_{j+1} < i < i+1 \leq r_j$, and then $m_i \geq m_{i+1}$.

Conversely, let be a collection of m_i verifying (i)–(ii). Then we define $w_j^1 \geq \dots \geq w_j^j$ as the conjugate partition of $m_{r_{j+1}+1} \geq \dots \geq m_{r_j}$ if $r_{j+1} < r_j$, and conditions (a) and (b) are obvious. ■

Finally, we remark that the numbers $\{m_i\}$ are the lengths of the Brunovsky chains of $\hat{f}: W \rightarrow X$. Hence, they are their Kronecker indices $\{\hat{k}_i\}$ up to permutation.

REFERENCES

- 1 N. Bourbaki, *Eléments de Mathématique*, Algèbre II, Hermann, Paris, 1964.
- 2 R. Bru, L. Rodman, and H. Schneider, Extensions of Jordan bases for invariant subspaces of a matrix, *Linear Algebra Appl.* 150:209–225 (1991).
- 3 I. Baragaña and I. Zaballa, Block similarity invariants of restrictions to (A, B) -invariant subspaces, *Linear Algebra Appl.* 220:31–62 (1995).
- 4 J. Ferrer and F. Puerta, Similarity of non-everywhere-defined linear maps, *Linear Algebra Appl.* 150:27–55 (1992).
- 5 J. Ferrer, F. Puerta, and X. Puerta, Geometric characterization and classification of marked subspaces, *Linear Algebra Appl.*, 235:15–34 (1996).
- 6 I. Gohberg, P. Lancaster, and L. Rodman, *Invariant Subspaces of Matrices with Applications*, Wiley, New York, 1986.

Received 15 February 1995; final manuscript accepted 6 November 1995